

On arc-disjoint Hamiltonian cycles in De Bruijn graphs

Z. Kása

Sapientia Hungarian University of Transylvania

email: kasa@ms.sapientia.ro

Presented at:

5th Joint Conference on Mathematics and Computer Science, Debrecen, June 9-12, 2004

and published in Hungarian:

De Bruijn-gráfok mint hálázati modellek (De Bruijn Graphs as Network Model), Műszaki Szemle/Technical Review, 43 (2008) 3-6

Abstract. We give two equivalent formulations of a conjecture [2,4] on the number of arc-disjoint Hamiltonian cycles in De Bruijn graphs.

A De Bruijn word of type (q, k) for a given q and k is a word over an alphabet with q letters, containing all k -length words exactly once. The length of such a word is $q^k + k - 1$. For example if $q = 3, k = 2$, then 0012202110 is a De Bruijn word of type $(3, 2)$.

For a q -letter alphabet $\mathcal{A}$ the De Bruijn graph $B(q, k)$ is defined as:

$B(q, k) = (V(q, k), E(q, k))$ with

- $V(q, k) = \mathcal{A}^k$ the set of vertices
- $E(q, k) = \mathcal{A}^{k+1}$ the set of directed arcs
- there is an arc from vertex $x_1x_2 \dots x_k$ to vertex $y_1y_2 \dots y_k$ if $x_2x_3 \dots x_k = y_1y_2 \dots y_{k-1}$ and this arc is denoted by $x_1x_2 \dots x_ky_k$.

In the De Bruijn graph $B(q, k)$ a path (i. e. a walk with distinct vertices) $a_1a_2 \dots a_k, a_2a_3 \dots a_{k+1}, \dots, a_{r-k+1}a_{r-k+2} \dots a_r, r > k$ corresponds to an r -length word $a_1a_2 \dots a_k a_{k+1} \dots a_r$, which is obtained by maximal overlapping of the neighboring vertices.

In $B(2, 3)$ the path 001, 010, 101 corresponds to the word 00101.

Every maximal length path in the graph $B(q, k)$ (which is a Hamiltonian one) corresponds to a De Bruijn word.

Algorithm to generate all De Bruijn words

$a_1, a_2, \dots, a_q$ the letters of an alphabet, with the values: $a_i = i - 1$ for $i = 1, 2, \dots, q$, S a vector to store the letters of a De Bruijn word, B a vector to store the states of words. $val(S_{i-k} \dots S_{i-1})$ represents the value of the number $S_{i-k} \dots S_{i-1}$ in base q .

Initially: $S_i := 0$ for all $i = 1, 2, \dots, k$, $B_0 := 1$ and $B_i := 0$ for other indices. The call of the procedure: DeBruijn ($S, k+1, k, B$).




Figure 1: De Bruijn graph $B(2, 2)$




Figure 2: De Bruijn graph $B(2, 3)$

```

procedure DeBruijn ( $S, i, k, B$ )
for  $j := 1$  to  $q$  do
   $S_i := a_j$ 
   $r := \text{val}(S_{i-k+1}S_{i-k+2} \dots S_i)$ 
  if  $B_r = 0$  then  $B_r := 1$ 
    DeBruijn( $S, i + 1, k, B$ )
     $B_r := 0$ 
  else if  $\text{length}(S) = q^k + k - 1$ 
    then write ( $S$ )
    exit for
  endif
endif
endfor
endprocedure

```

In the directed graph $B(q, k)$ there always exists an Eulerian circuit because it is connected and all its vertices have the same indegree and outdegree q . An Eulerian circuit in $B(q, k)$ is a Hamiltonian path in $B(q, k + 1)$ (which always can be continued in a Hamiltonian cycle).




Figure 3: De Bruijn graph $B(3, 2)$

Conjecture 1. [2, 4]

In the De Bruijn graph $B(q, k)$ for $q \geq 2$ and $k > 1$ the number of arc-disjoint Hamiltonian cycles is $q - 1$.

Let us define a morphism μ on words over an alphabet $A = \{0, 1, \dots, q - 1\}$:

$$\begin{aligned} \mu(0) &= 0 \\ \mu(i) &= i + 1, \quad \text{if } 1 \leq i < q - 1 \\ \mu(q - 1) &= 1 \end{aligned}$$

It is easy to see that $\mu^{q-1}(u) = u$ for any $u \in A^*$.

Example:

Let be a word $w = 00102113230331220$ on the alphabet $A = \{0, 1, 2, 3\}$.

$$\mu(w) = 00203221310112330$$

$$\mu^2(w) = 00301332120223110$$

From a De Bruijn word we obtain a Hamiltonian cycle. Let an H_0 be such a Hamiltonian cycle. Using μ^k for $k = 1, 2, \dots, q - 2$ we will obtain De Bruijn words corresponding to the Hamiltonian cycles $H_1, H_2, \dots, H_{q-2}$.

Conjecture 2.

In the De Bruijn graph $B(q, k)$ for $q > 2, k > 1$ there exists a Hamiltonian cycle H_0 such that the Hamiltonian cycles $H_1, H_2, \dots, H_{q-2}$ (obtained from H_0 by using the morphisms $\mu_k, k = 1, 2, \dots, q-2$), together with H_0 are arc-disjoint Hamiltonian cycles.

$B(3, 2)$

$H_0 : 0011220210,$

$H_1 : 0022110120$

$B(3, 3)$

$H_0 : 00010021011022202012111221200,$

$H_1 : 00020012022011101021222112100$

$B(4, 2)$

$H_0 : 00102113230331220,$

$H_1 : 00203221310112330,$

$H_2 : 00301332120223110$

$B(5, 2)$

$H_0 : 00102112041422430332313440,$

$H_1 : 00203223012133140443424110,$

$H_2 : 00304334023244210114131220,$

$H_3 : 00401441034311320221242330$

Two words u and v are equivalent if $v = \mu^k(u)$ for some $1 \leq k \leq q-2$

If $v = \mu^k(u)$, $1 \leq k \leq q-2$, then $u = \mu^{q-1-k}(v)$ because, applying μ^{q-1-k} to $v = \mu^k(u)$ we obtain $\mu^{q-1-k}(v) = \mu^{q-1-k}(\mu^k(u)) = \mu^{q-1}(u) = u$.

In the set of all De Bruijn words over an alphabet $A = \{0, 1, \dots, q-1\}$ this equivalence relation will introduce a partition of De Bruijn words in equivalent classes.

An equivalent assertion to Conjecture 2.

In the De Bruijn graph $B(q, k)$ for $q > 2, k > 1$ there exist the Hamiltonian cycles $H_0, H_1, H_2, \dots, H_{q-2}$ such that they correspond to De Bruijn words $B_0, B_1, B_2, \dots, B_{q-2}$ from the same equivalence class.




Figure 4: Arc-disjoint Hamiltonian cycles in De Bruijn graph $B(3, 2)$

References

1. M. C. Anisiu, Z. Blázsik, Z. Kása, *Maximal complexity of finite words*, Pure Math. and Appl. Vol. **13** (2002) No. 1–2, pp. 39–48.
2. J. Bond, A. Iványi, *Modelling of interconnection networks using De Bruijn graphs*, Third Conference of Program Designer, Ed. A. Iványi, Budapest, 1987, 75–87.
3. N. G. de Bruijn, *A combinatorial problem*, Nederl. Akad. Wetensch. Proc. 49 (1946), 758–764.
4. Sophie Gire, *Réseaux d'interconnexion*, Option Maîtrise Informatique, 1996–97,
http://fastnet.univ-brest.fr/~gire/COURS/OPTION_RESEAUX/node48.html
5. A. Iványi, M. Horváth, *Perfect sequences*, ICAI'2004, Eger, January 28, 2004.
6. M. Lothaire, *Combinatorics on words*, Addison-Wesley, Reading, MA, 1983.
7. M. H. Martin, *A problem in arrangements*, Bull. A.M.S. 40 (1934), 859–864.